

Dr. Adhershitha AR

Assistant Professor, Department of Prosthodontics, Government Dental College, Kottayam.

- Reason for brittleness of endodontically treated tooth
 - Micro cracks in remaining tooth structure produced by endodontic procedures.
 - Weakened collagen intermolecular cross-links in Dentine
 - Loss of internal & external tooth structure
 - All the above
- Average number of deglutitions per day
 - 450
 - 500
 - 550
 - 600
- The most effective way to resolve inflammation of oral tissues is to keep the denture out of the mouth for at least ____ hours before impressions are made.
 - 12
 - 18
 - 24
 - 48
- Chewing centre is located in
 - Pons
 - Thalamus
 - Frontal lobe
 - Cortex
- Genial tubercle becomes prominent in
 - Class I patients
 - Class II patients
 - Class III patients
 - In resorbed ridge
- Beyrton's point is
 - 13 mm in front of anterior margin of meatus
 - 13 mm from posterior margin of tragus to canthus
 - 13mm from foot of tragus to canthus
 - 10 mm anterior to centre of external auditory meatus and 7mm below Frankfort plane
- Path of insertion for a $\frac{3}{4}$ crown
 - Parallel to occlusal 1/3
 - Parallel to occlusal 2/3
 - Parallel to cervical 1/3
 - Parallel to long axis
- Functional cusp bevel gives
 - Help to preserve periodontium
 - Provides resistance form
 - Provides structural durability
 - Help to preserve tooth structure
- A parallel pin added to prefabricated resin post for antirotational stabilization and some additional retention is called
 - Copost
 - Parapost
 - Auxillary post
 - Additional post
- Arc-fixed partial denture is otherwise called as
 - Working denture
 - Functional denture
 - Interim denture
 - Perel pontic

1. D, 2. B, 3. C, 4. A, 5. D, 6. B, 7. D, 8. C, 9. B, 10. D

Answers